



## Recombinant CD200 Receptor 1 (CD200R1)

Catalog No.: TP01488

50µg

### Sequence Information

**Species:** Mouse

**Gene ID:**57781

**Swiss Prot:**Q9ES57

**Synonyms:**OX2R; HCRT2; CD200R;

MOX2R; MOX2 Receptor; Cell surface glycoprotein CD200 receptor 1; CD200 cell surface glycoprotein receptor; Cell surface glycoprotein OX2 receptor 1

**Residues:**Thr26~Pro238

TDKNQTTQNNSSSPLTQVNTTVSVQIGTKALLCCFSIPLTKAVLITWIIKLRGL  
PSCTIAYKVDTKTNETSCLGRNITWASTPDHSPQLQISAVTLQHEGTYTCETVT  
PEGNFEKNYDLQVLVPPEVTYFPEKNRSVCEAMAGKPAAQISWSPDGDVTTTS  
ESHNGTVTVRSTCHWEQNNVSDVSCIVSHLTGNQSLSELSRGGNQSLRP

### Product Information

**Source:** Prokaryotic expression.

**Host:** *E.coli*

**Tags:** N-terminal His-Tag

**Subcellular Location:** Membrane.

**Purity:** >90%

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

**Original Concentration:** 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 6.0

**Predicted Molecular Mass:** 26.7kDa

**Accurate Molecular Mass:** 33kDa as determined by SDS-PAGE reducing conditions.

### [ USAGE ]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE



### **[ IMPORTANT NOTE ]**

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.