



Recombinant Human Estrogen Receptor Beta (ERb)

Catalog No.: TP09318

100µg

Sequence Information

Species: Human

Gene ID:2100

Swiss Prot:Q92731

Synonyms: ESR2

Residues:Met46-Gln530

MTFYSPAVMNYSIPSNVTNLEGGPGRQTTSPNVLWPTPGHLSPLVHRQLSHLYAEP
QKSPWCEARSLEHTLPVNRETLKRKVSNGRCASPVTPGSGSKRDAHFCAVCSFYASG
YHYGVWSCEGCKAFFKRSIQGHNDYICPATNQCTIDKNRRKSCQACRLRKCYEVGM
VKCGSRRERCGYRLVRRQRSADQLHCAGKAKRSGGHAPRVRELLLDALSPEQLVL
TLLEAEPHPVLISRPSAPFTEASMMMSLTKLADKELVHMISWAKKIPGFVLSLFDQVR
LLESCWMEVLMMGLMWRSIDHPGKLIFAPDLVDRDEGKCVGEILEIFDMLLATTSRF
RELKLQHKEYLCVKAMILLNSSMYPLVTATQDADSSRKLHLLNAVTDALVWVIAKSGI
SSQQQSMRLANLLMLLSHVRHASNKGMEHLLNMKCKNVVPVYDLLLEMLNAHVLRG
CKSSITGSECSPAEDSKSKEGSQNPQSQ

Product Information

Source: Prokaryotic expression.

Host: *E. coli*

Tags: N-terminal His-Tag

Subcellular Location: Membrane.

Purity: >80%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 55kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at



37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.