



## Recombinant Sepiapterin Reductase (SPR)

Catalog No.: TP03886

100µg

### Sequence Information

**Species:** Human

**Gene ID:**6697

**Swiss Prot:**P35270

**Synonyms:**SDR38C1;

7,8-Dihydrobiopterin:NADP+

Oxidoreductase; Short Chain

Dehydrogenase/Reductase Family

38C,Member 1

**Residues:**Met1~Lys261

MEGGLGRAVCLLTGASRGFGRTLAPLLASLLSPGSVLVLSARNDEALRQLEAEL

GAERSGLRVVRVPADLGAEAGLQQLLGALRELPRPKGLQRLLLINNAGSLGDVS

KGFVDLS DSTQVNNY WALNLT SMLCLTSSVLKAFD SPGLNRTVVNISSLCALQ

PFKGWALYCAGKAARDMLFQVLALEEPNVRVLNYAPGPLD TDMQQLARETSVDP

DMRKG LQELKAKGKLVDCKVSAQKLLSLLEKDEFKSGAHVDFYDK

### Product Information

**Source:** Recombinant expression.

**Host:** *E.coli*

**Tags:** Two N-terminal Tags, His-tag and GST-tag

**Subcellular Location:** Cytoplasm.

**Purity:** >90%

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

**Original Concentration:** 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 8.3

**Predicted Molecular Mass:** 58.0kDa

**Accurate Molecular Mass:** 58kDa as determined by SDS-PAGE reducing conditions.

### [ USAGE ]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]


**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



**Figure 1. SDS-PAGE**

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.