

## **Recombinant Meningioma Expressed Antigen 5 (MGEA5)**

Catalog No.: TP09526 100µg

**Sequence Information** 

Species: Human Gene ID:10724

Swiss Prot: O60502 Synonyms: NAGase; MEA5; NCOAT;

Hexosaminidase C;

N-Acetylhexosaminidase; Hyaluronidase;

Meningioma-Expressed Antigen 5;

Nuclear Cytoplasmic O-GlcNAcase And

Acetyltransferase

Residues: Ser6~Lys240

SOATLEERESELSSNPAASAGASLEPPAAPAPGEDNPAGAGGAAVAGAAGGARR

FLCGVVEGFYGRPWVMEQRKELFRRLQKWELNTYLYAPKDDYKHRMFWREMYSV

EEAEQLMTLISAAREYEIEFIYAISPGLDITFSNPKEVSTLKRKLDQVSQFGCR

SFALLFDDIDHNMCAADKEVFSSFAHAQVSITNEIYQYLGEPETFLFCPTEYCG

**TFCYPNVSQSPYLRTVGEK** 

**Product Information** 

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His-Tag

**Subcellular Location:** Nucleus, Cytoplasm, Chromosome.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.2

Predicted Molecular Mass: 27.5kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]



Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE

## [IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.