

Eukaryotic Cluster Of Differentiation 14 (CD14)

Catalog No.: TP09572

50µg

Sequence Information

Species: Human

Gene ID: 929

Swiss Prot: P08571

Synonyms: Monocyte differentiation antigen

CD14; Myeloid cell-specific leucine-rich
glycoprotein

Residues: Thr20-Cys352

TTPEPCELDDEDFRCVCNFSEPQPDWSEAFQCVSAVEVEIHAGGLNLEPFLKRV
DADADPRQYADTVKALRVRRLLTVGAAQVPAQLLVGALRVLAYSRKELTLEDLK
ITGTMPPPLPLEATGLALSSLRLRNVSATGRSWLAELQQWLKPKLVLSIAQAH
SPAFSCEQVRAFPALTSLDLSDNPGLGERGLMAALCPHKFPAIQNLALRNTGME
TPTGVCAALAAAGVQPHSLDLSHNSLRATVNPSAPRCMWSSALNSLNSFAGLE
QVPKGLPAKLRVLDLSCNRLNRAPQPDELPEVDNLTLDGNPFLVPGTALPHEGS
MNSGVVPAC

Product Information

Source: Eukaryotic expression.

Host: 293F cell

Tags: C-terminal His Tag

Subcellular Location: Membrane, Secreted.

Purity: >97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.4

Predicted Molecular Mass: 54.6kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.