

## Recombinant Human CD300 Antigen Like Family Member C (CD300c)

Catalog No.: TP05474

50µg

### Sequence Information

**Species:** Human

**Swiss Prot:**Q08708

**Gene ID:**

**Synonyms:**CMRF35-Like Molecule 6; CLM-6;  
CD300 Antigen-Like Family Member C;  
CMRF35-A1; CMRF-35;  
Immunoglobulin Superfamily Member  
16; IgSF16; CD300c; CD300C;  
CMRF35; CMRF35A; CMRF35A1;  
IGSF16

**Residues:**Gly21-Arg183

GYFPLSHPMTVAGPVGGSLSVQCRYEKEHRTL NKFWCRPPQILRCDKIVETKGS  
AGKRNGRVSIIRDSPANLSFTVTLENL TEEDAGTYWCGVDTPWLRDFHDPIVEVE  
VSVFPAGTTTASSPQSSMGTS GPPTKLPVHTWPSVTRKDSPEPSHPGSLFSNV  
R

### Product Information

**Source:** Eukaryotic expression.

**Host:** 293F cell

**Tags:** C-terminal His-Tag

**Subcellular Location:** Cell membrane.

**Purity:** >95%

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

**Original Concentration:** 1000µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 8.0

**Predicted Molecular Mass:** 40kDa

**Accurate Molecular Mass:** 40kDa as determined by SDS-PAGE reducing conditions.

### [ USAGE ]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.