



Recombinant Human DDX56 Protein

Catalog No.: TP01855 100µg

Sequence Information

Species: Human

Gene ID:54606

Swiss Prot:Q9NY93

Synonyms: DDX21, DDX26, NOH61

Residues:Met1-Ser547

MEDSEALGFEHMGGLDPRLLQAVTDLGWSRPTLIQEKAIPALLEGKDLLARARTGSGKTAAYAI PMLQLLLHRKATGP
VVEQAVRGLVLVPTKELARQAQSMIQQLATYCARDVRVANVSAAEDSVSQRAVLMEKPDVVVGTPSRILSHLQQDSL
KLRDSLELLLVDEADLLFSFGFEEELKSL LCHLPRIYQAF LMSATFNEDVQALKELILHNPVTLKLQESQLPGPDQL
QQFQVVCETEEDKFLLLYALLKLSLIRGKSL LFVNTLERSYRLRLFLEQFSIPTCVLNGELPLRSRCHIIISQFNQGF
YDCVIATDAEVLGAPVKGKRRGRGPKGDKASDPEAGVARGIDFHHVSAVLNFDLPPTPEAYIHRAGRTARANNPGIV
LTFVLPTEQFHLGKIEELLSGENRGPI LLYPYQFRMEEIEGFRYRCRDAMRSVTKQAI REARLKEIKEELLHSEKLKT
YFEDNPRDLQLLRHDLPLHPAVVKPHLGHPDYLVPPALRGLVRPHKKRKKLSSSCRKAKRAKSQNPLRSFKHKGKK
FRPTAKPS

Product Information

Source: Prokaryotic expression.

Host: *E. coli*

Tags: N-terminal His-Tag

Subcellular Location: Secreted.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS (PH7.4)) , containing 5% Trehalose.

Original Concentration: 400µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.4

Predicted Molecular Mass: 64kDa

Accurate Molecular Mass: 64kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in PBS (pH7.4) to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure . SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.