

### Recombinant Isocitrate dehydrogenase [NADP], mitochondrial (IDHP)

Catalog No.: TP10059 100µg

#### **Sequence Information**

Species: Mouse Gene ID:269951

Swiss Prot:P54071 Synonyms:Idh2

Residues: Thr 40-Gln 452

TWAPAALTVPSWPEQPRRHYAEKRIKVEKPVVEMDGDEMTRIIWQFIKEKLILP

HVDVQLKYFDLGLPNRDQTNDQVTIDSALATQKYSVAVKCATITPDEARVEEFK

LKKMWKSPNGTIRNILGGTVFREPIICKNIPRLVPGWTKPITIGRHAHGDQYKA

TDFVVDRAGTFKLVFTPKDGSSAKEWEVYNFPAGGVGMGMYNTDESISGFAHSC

FQYSIQKKWPLYLSTKNTILKAYDGRFKDIFQEIFDKHYKTDFDKNKIWYEHRL

 ${\tt IDDMVAQVLKSSGGFVWACKNYDGDVQSDILAQGFGSLGLMTSVLVCPDGKTIE}$ 

AEAAHGTVTRHYREHQKGRPTSTNPIASIFAWTRGLEHRGKLDGNQDLIRFAQT

LEKVCVQTVESGAMTKDLAGCIHGLSNVKLNEHFLNTTDFLDTIKSNLDRALGK

Q

#### **Product Information**

**Source:** Recombinant expression.

Host: E.coli

**Tags:** N-terminal His-Tag

Subcellular Location: Mitochondrion .

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 55.3kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing conditions.

## [USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.



### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



Figure 1. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.