



Recombinant Human α -Synuclein (SNCA) Protein (Active)

Catalog No.: TP10117 100 μ g

Sequence Information

Species: Human

Gene ID:6622

Swiss Prot:P37840

Synonyms:SNCA, NACP, PARK1, alpha-Synuclein

Residues:Met1-Ala140

MDVFMKGLSKAKEGVVAAAEKTKQGVAEAAGKTKEGVLYVGSKTKEGVVHGVATVAEKTKEQVTNVGGAVVTGVTAVAQKTVEGAGSIAAATGF
VKKDQLGKNEEGAPQEGILEDMPVDPDNEAYEMPSEEGYQDYEP

Product Information

Source: Recombinant expression.

Host: *E. coli*

Tags: N-terminal His-Tag

Subcellular Location: Membrane, Cytoplasm, Golgi apparatus.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200 μ g/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.7

Predicted Molecular Mass: 14.4kDa

Accurate Molecular Mass:14.4kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[ACTIVITY]

100 μ M alpha synuclein protein monomer , seeded with 10 μ M alpha synuclein protein aggregate in 25 μ M Thioflavin T (ab120751) (PBS pH 7.4, 100 μ l reaction volume), generated a fluorescence intensity of 13,000 Relative Fluorescence Units after incubation at 37° C with shaking at 600 rpm for 24 hours

[IDENTIFICATION]




Figure . SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.