

Recombinant Human Programmed cell death protein 4(PDCD4)

Catalog No.: **TP10464** 100µg

Sequence Information

Species: Human Swiss Prot:Q53EL6

Gene ID:27250

Synonyms:Programmed cell death 4 isoform 1,

Neoplastic transformation inhibitor,

H731.

Residues:Met1-Asn325

MDVENEQILNVNPADPDNLSDSLFSGDEENAGTEEIKNEINGNWISASSINEAR INAKAKRRLRKNSSRDSGRGDSVSDSGSDALRSGLTVPTSPKGRLLDRRSRSGK GRGLPKKGGAGGKGVWGTPGQVYDVEEVDVKDPNYDDDQENCVYETVVLPLDER AFEKTLTPIIQEYFEHGDTNEVAEMLRDLNLGEMKSGVPVLAVSLALEGKASHR EMTSKLLSDLCGTVMSTTDVEKSFDKLLKDLPELALDTPRAPQLVGQFIARAVG DGILCNTYIDSYKGTVDCVQARAALDKATVLLSMSKGGKRKDSVWGSGGGQQSV N

Product Information

Source: Recombinant expression. Host: *E.coli* Tags: N-terminal His-Tag Subcellular Location: Cytoplasm. Purity: >90% Traits: Freeze-dried powder Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 4.7 Predicted Molecular Mass: 38.4kDa Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]

KDa		
180 130	and the second se	
130		
70		
	-	
55		
40		
		_
_		
35		
25		
20		
_		
_		
_		
15		
15		
_		
_		
_		
10		

Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.