

## Recombinant Immunoglobulin A2 (IgA2)

Catalog No.: TP10481 100µg

**Sequence Information** 

Species: Human Gene ID:

Swiss Prot:A0A0G2JMB2 Synonyms:IGHA2; Immunoglobulin

Heavy Constant Alpha 2; A2m Marker

Residues: Ser1-Tyr339

 ${\tt SPTSPKVFPLSLDSTPQDGNVVVACLVQGFFPQEPLSVTWSESGQNVTARNF}$ 

PPSQDASGDLYTTSSQLTLPATQCPDGKSVTCHVKHYTNPSQDVTVPCPVPP

PPPCCHPRLSLHRPALEDLLLGSEANLTCTLTGLRDASGATFTWTPSSGKSAV

QGPPERDLCGCYSVSSVLPGCAQPWNHGETFTCTAAHPELKTPLTANITKSG

NTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDVLVRWLQGSQELPREK

YLTWASRQEPSQGTTTFAVTSILRVAAEDWKKGDTFSCHVGHEALPLAFTQ

KTIDRLAGKPTHVNVSVVIAEVDGTCY

**Product Information** 

**Source:** Recombinant expression.

Host: E.coli

Tags: Two N-terminal Tags, His-tag and SUMO-tag

Subcellular Location: Membrane, Secreted.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.4

Predicted Molecular Mass: 56.0kDa

Accurate Molecular Mass: 56kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[ STORAGE AND STABILITY ]



Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.