

## Recombinant Immunoglobulin E (IgE)

Catalog No.: TP11444 100µg

**Sequence Information** 

Species: Mouse Gene ID:

Swiss Prot: P06336 Synonyms: IGHE; Immunoglobulin Heavy

Constant Epsilon; Ig epsilon chain C

region

Residues: Ser1-Ser421

SIRNPQLYPLKPCKGTASMTLGCLVKDYFPNPVTVTWYSDSLNMSTVNFPALGS

ELKVTTSQVTSWGKSAKNFTCHVTHPPSFNESRTILVRPVNITEPTLELLHSSC

DPNAFHSTIQLYCFIYGHILNDVSVSWLMDDREITDTLAQTVLIKEEGKLASTC

 ${\tt SKLNITEQQWMSESTFTCKVTSQGVDYLAHTRRCPDHEPRGVITYLIPPSPLDL}$ 

YQNGAPKLTCLVVDLESEKNVNVTWNQEKKTSVSASQWYTKHHNNATTSITSIL

PVVAKDWIEGYGYQCIVDHPDFPKPIVRSITKTPGQRSAPEVYVFPPPEEESED

KRTLTCLIQNFFPEDISVQWLGDGKLISNSQHSTTTPLKSNGSNQGFFIFSRLE

VAKTLWTQRKQFTCQVIHEALQKPRKLEKTISTSLGNTSLRPS

**Product Information** 

**Source:** Recombinant expression.

Host: E.coli

Tags: N-terminal His Tag

Subcellular Location: Secreted.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.1

Predicted Molecular Mass: 51.0kDa

Accurate Molecular Mass: 51kDa as determined by SDS-PAGE reducing conditions.

[USAGE]



Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.