

### Recombinant Transcription factor AP-2 gamma (TFAP2C)

Catalog No.: TP10506 100µg

#### **Sequence Information**

Species: Human Gene ID:7022

Swiss Prot:Q92754 Synonyms:AP2C

Residues: Met1-Lys450

MLWKITDNVKYEEDCEDRHDGSSNGNPRVPHLSSAGQHLYSPAPPLSHTGVAEY

QPPPYFPPPYQQLAYSQSADPYSHLGEAYAAAINPLHQPAPTGSQQQAWPGRQS

QEGAGLPSHHGRPAGLLPHLSGLEAGAVSARRDAYRRSDLLLPHAHALDAAGLA

 ${\tt ENLGLHDMPHQMDEVQNVDDQHLLLHDQTVIRKGPISMTKNPLNLPCQKELVGA}$ 

VMNPTEVFCSVPGRLSLLSSTSKYKVTVAEVQRRLSPPECLNASLLGGVLRRAK

SKNGGRSLREKLDKIGLNLPAGRRKAAHVTLLTSLVEGEAVHLARDFAYVCEAE

FPSKPVAEYLTRPHLGGRNEMAARKNMLLAAQQLCKEFTELLSQDRTPHGTSRL

APVLETNIQNCLSHFSLITHGFGSQAICAAVSALQNYIKEALIVIDKSYMNPGD

**QSPADSNKTLEKMEKHRK** 

#### **Product Information**

**Source:** Recombinant expression.

Host: E.coli

Tags: N-terminal His and KSI Tag
Subcellular Location: Nucleus.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.9

Predicted Molecular Mass: 65.2kDa

Accurate Molecular Mass: 65kDa as determined by SDS-PAGE reducing conditions.

# [USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.



### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]




Figure 1. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.