

## Recombinant Phosphoinositide-3-Kinase Class 3 (PIK3C3)

Catalog No.: TP11652 100µg

**Sequence Information** 

Species: Human Gene ID:5289

Swiss Prot:Q8NEB9 Synonyms:Vps34; Phosphatidylinositol

3-kinase p100 subunit;

Phosphatidylinositol 3-kinase catalytic

subunit type 3

Residues: Tyr631-Trp885

YPVIFKHGDDLRQDQLILQIISLMDKLLRKENLDLKLTPYKVLATSTKHGFMQF

IQSVPVAEVLDTEGSIQNFFRKYAPSENGPNGISAEVMDTYVKSCAGYCVITYI

LGVGDRHLDNLLLTKTGKLFHIDFGYILGRDPKPLPPPMKLNKEMVEGMGGTQS

EQYQEFRKQCYTAFLHLRRYSNLILNLFSLMVDANIPDIALEPDKTVKKVQDKF

RLDLSDEEAVHYMQSLIDESVHALFAAVVEQIHKFAQYW

**Product Information** 

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His Tag

Subcellular Location: Late endosome.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.5

Predicted Molecular Mass: 32.9kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[ STORAGE AND STABILITY ]



Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.