

Recombinant Human Coagulation Factor VII / FVII / F7 Protein

Catalog No.: TP07445

50µg

Sequence Information

Species: Human

Gene ID:2155

Swiss Prot:P08709-2

Synonyms: Coagulation factor 7;Coagulation factor VII;SPCA

Residues:Met1-Pro444

MVSQALRLLCLLLGLQGCLAAVFVTQEEAHGVLHRRRRANAFLEELRPGSLERE
CKEEQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYIC
FCLPAFEGRNCETHKDDQLICVNEGGCEQYCSHTGTRSCRCHEGYSLLADG
VSCTPTVEYPCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLC
GGTLINTIWWVSAAHCFDKIKNWRNLI AVLGEHDLSEHDGDEQSRRVAQVIIPS
TYVPGTTNHDIALRLHQPVVLT DHVVPLCLPERTFSERTLAFVRFSLVSGWGQ
LLDRGATALELMVLNVPRLMTQDCLQQSRKVG DSPNITEYMF CAGYSDGSKDSC
KGDSGGPHATHYRG TWYLTGIVSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSE
PRPGVLLRAPFP

Product Information

Source: Eukaryotic expression.

Host: 293F cell

Tags: C-terminal His-Tag

Subcellular Location: Secreted .

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 1000µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.9

Predicted Molecular Mass: 52.7kDa

Accurate Molecular Mass: 53kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.